arXiv:1112.3754v1 [quant-ph] 16 Dec 2011

Geometrical structures of multipartite quantum systems

Hoshang Heydari
Physics Department, Stockholm university 10691 Stockholm Sweden

Email: hoshang@fysik.su.se

December 19, 2011

Abstract

In this paper I will investigate geometrical structures of multipartite quantum systems based on complex projective varieties. These varieties are important in characterization of quantum entangled states. In particular I will establish relation between multi-projective Segre varieties and multip-qubit quantum states. I also will discuss other geometrical approaches such as toric varieties to visualize complex multipartite quantum systems.

1 Introduction

Characterization of multipartite quantum systems is very interesting research topic in the foundations of quantum theory and has many applications in the field quantum information and quantum computing. And in particular geometrical structures of multipartite quantum entangled pure states are of special importance. In this paper we will review the construction of Segre variety for multi-qubit states. We will also show a construction of geometrical measure of entanglement based on the Segre variety for multi-qubit systems. Finally we will establish a relation between the Segre variety, toric variety, and multi-qubit quantum systems. The relation could be used as a tool to visualize entanglement properties of multi-qubit states. Let $\mathcal{Q}_j,\ j=1,2,\ldots m$ be quantum systems with underlying Hilbert spaces $\mathcal{H}_{\mathcal{Q}_j}$. Then the Hilbert space of a multi-qubit systems $\mathcal{Q}$, is given by $\mathcal{H}_{\mathcal{Q}}=\mathcal{H}_{\mathcal{Q}_m}\otimes\mathcal{H}_{\mathcal{Q}_{m-1}}\otimes\cdots\otimes\mathcal{H}_{\mathcal{Q}_1}$, where $\mathcal{H}_{\mathcal{Q}_j}=\mathbf{C}^2$ and $\dim\mathcal{H}_{\mathcal{Q}}=2^m$. Now, let

$$|\Psi\rangle=\sum_{x_m=0}^{1}\sum_{x_{m-1}=0}^{1}\cdots\sum_{x_1=0}^{1}\alpha_{x_m x_{m-1}\cdots x_1}|x_m x_{m-1}\cdots x_1\rangle, \tag{1}$$

be a vector in $\mathcal{H}_{\mathcal{Q}}$, where $|x_m x_{m-1}\cdots\ x_1\rangle=|x_m\rangle\otimes|x_{m-1}\rangle\otimes\cdots\otimes|x_1\rangle$ are orthonormal basis in $\mathcal{H}_{\mathcal{Q}}$ and $\alpha_{x_m x_{m-1}\cdots x_1}\in\mathbf{C}$. Then the quantum states are normalized vectors in $\mathcal{P}(\mathcal{H}_{\mathcal{Q}})=\mathcal{H}_{\mathcal{Q}}/\sim$. Moreover, let $\rho_{\mathcal{Q}}=\sum_{i=1}^{\mathrm{N}}p_i|\Psi_i\rangle\langle\Psi_i|$, for all $0\leq p_i\leq 1$ and $\sum_{i=1}^{\mathrm{N}}p_i=1$, denotes a density operator acting on the Hilbert space $\mathcal{H}_{\mathcal{Q}}$.

The density operator $\rho_{\mathcal{Q}}$ is said to be fully separable, which we will denote by $\rho_{\mathcal{Q}}^{sep}$, with respect to the Hilbert space decomposition, if it can be written as $\rho_{\mathcal{Q}}^{sep} = \sum_{i=1}^{\mathrm{N}} p_i \bigotimes_{j=1}^{m} \rho_{\mathcal{Q}_j}^{i}$, where $\rho_{\mathcal{Q}_j}^{i}$ denotes a density operator on Hilbert space $\mathcal{H}_{\mathcal{Q}_j}$. If $\rho_{\mathcal{Q}}^{p}$ represents a pure state, then the quantum system is fully separable if $\rho_{\mathcal{Q}}^{p}$ can be written as $\rho_{\mathcal{Q}}^{sep} = \bigotimes_{j=1}^{m} \rho_{\mathcal{Q}_j}$, where $\rho_{\mathcal{Q}_j}$ is the density operator on $\mathcal{H}_{\mathcal{Q}_j}$. If a state is not separable, then it is said to be an entangled state.

2 Projective geometry

In this section we give a short introduction to variety. Let $\mathbf{C}[z] = \mathbf{C}[z_1, z_2, \dots, z_n]$ denotes the polynomial algebra in n variables with complex coefficients. Then, given a set of r polynomials $\{g_1, g_2, \dots, g_r\}$ with $g_i \in \mathbf{C}[z]$, we define a complex affine variety as

$$\mathcal{V}_{\mathbf{C}}(g_1, g_2, \dots, g_r) = \{P \in \mathbf{C}^n : g_i(P) = 0 \ \forall \ 1 \leq i \leq r\}, \tag{2}$$

where $P \in \mathbf{C}^n$ is called a point of $\mathbf{C}^n$ and if $P = (a_1, a_2, \dots, a_n)$ with $a_j \in \mathbf{C}$, then a_j is called the coordinates of P. A complex projective space $\mathbf{CP}^n$ is defined to be the set of lines through the origin in $\mathbf{C}^{n+1}$, that is,

$$\mathbf{CP}^n = \frac{\mathbf{C}^{n+1} - \{0\}}{u \sim v}, \ \lambda \in \mathbf{C} - 0, \ v_i = \lambda u_i \ \forall \ 0 \leq i \leq n+1, \tag{3}$$

where $u = (u_1, \dots, u_{n+1})$ and $v = (v_1, \dots, v_{n+1})$. Given a set of homogeneous polynomials $\{g_1, g_2, \dots, g_r\}$ with $g_i \in \mathbf{C}[z]$, we define a complex projective variety as

$$\mathcal{V}(g_1, \dots, g_r) = \{O \in \mathbf{CP}^n : g_i(O) = 0 \ \forall \ 1 \leq i \leq r\}, \tag{4}$$

where $O = [a_1, a_2, \dots, a_{n+1}]$ denotes the equivalent class of point $\{\alpha_1, \alpha_2, \dots, \alpha_{n+1}\} \in \mathbf{C}^{n+1}$. We can view the affine complex variety $\mathcal{V}_{\mathbf{C}}(g_1, g_2, \dots, g_r) \subset \mathbf{C}^{n+1}$ as a complex cone over the complex projective variety $\mathcal{V}(g_1, g_2, \dots, g_r)$.

We can map the product of spaces $\underbrace{\mathbf{CP}^1 \times \mathbf{CP}^1 \times \cdots \times \mathbf{CP}^1}_{m \text{ times}}$ into a projective space by its Segre embedding as follows. The Segre map is given by

$$\mathcal{S}_{2,\dots,2} : \mathbf{CP}^1 \times \mathbf{CP}^1 \times \cdots \times \mathbf{CP}^1 \quad \longrightarrow \quad \mathbf{CP}^{2^m - 1}, \tag{5}$$

is defined by $((\alpha_0^1, \alpha_1^1), \dots, (\alpha_0^m, \alpha_1^m)) \longmapsto (\alpha_{i_m}^m \alpha_{i_{m-1}}^{m-1} \cdots \alpha_{i_1}^1)$, where (α_0^i, α_1^i) is points defined on the ith complex projective space $\mathbf{CP}^1$ and $\alpha_{i_m i_{m-1} \cdots i_1}$,$0 \leq i_s \leq 1$ be a homogeneous coordinate-function on $\mathbf{CP}^{2^m - 1}$. Moreover, let us consider a multi-qubit quantum system and let $\mathcal{A} = \left(\alpha_{i_m i_{m-1} \dots i_1}\right)_{0 \leq i_s \leq 1}$, for all $j = 1, 2, \dots, m$. $\mathcal{A}$ can be realized as the following set $\{(i_1, i_2, \dots, i_m) : 1 \leq i_s \leq 2, \forall \ s\}$, in which each point $(i_m, i_{m-1}, \dots, i_1)$ is assigned the value $\alpha_{i_m i_{m-1} \dots i_1}$. For each $s = 1, 2, \dots, m$, a two-by-two minor about the j-th coordinate of $\mathcal{A}$ is given by

$$\mathcal{I}_{\mathcal{A}}^{m} = \alpha_{x_m x_{m-1} \dots x_1} \alpha_{y_m y_{m-1} \dots y_1} - \alpha_{x_m x_{m-1} \dots x_{s+1} y_s x_{s-1} \dots x_1} \alpha_{y_m y_{m-1} \dots y_{s+1} x_s y_{s-1} \dots y_m} \tag{6}$$

Then the ideal $\mathcal{I}^m_{\mathcal{A}}$ is generated by The image of the Segre embedding $\mathrm{Im}(\mathcal{S}_{2,2,\dots,2})$, which again is an intersection of families of quadric hypersurfaces in $\mathbf{CP}^{2^m-1}$, is called Segre variety and it is given by

$$\mathrm{Im}(\mathcal{S}_{2,2,\dots,2}) \quad = \quad \bigcap_{\forall s} \mathcal{V}\left(\mathcal{I}^m_{\mathcal{A}}\right). \tag{7}$$

This is the space of separable multi-qubit states. Moreover, we propose a measure of entanglement for general pure multipartite states based on modified Segre variety as follows

$$\begin{aligned}\mathcal{F}(\mathcal{Q}^p_m(2,2\dots,2)) = (\mathcal{N} \sum_{\forall\sigma\in\mathrm{Perm}(u)} \sum_{k_j,l_j,j=1,2,\dots,m} \\ |\alpha_{k_1k_2\dots k_m}\alpha_{l_1l_2\dots l_m} - \alpha_{\sigma(k_1)\sigma(k_2)\dots\sigma(k_m)}\alpha_{\sigma(l_1)\sigma(l_2)\dots\sigma(l_m)}|^2)^{\frac{1}{2}},\end{aligned} \tag{8}$$

where $\sigma \in \mathrm{Perm}(u)$ denotes all possible sets of permutations of indices for which $k_1k_2\dots k_m$ are replace by $l_1l_2\dots l_m$, and u is the number of indices to permute.

As an example we will discuss the four-qubit state in which we first encounter these new varieties. For this quantum system we can partition the Segre embedding as follows:

$$\begin{tikzcd}
\mathbf{P}^1\times\mathbf{P}^1\times\mathbf{P}^1\times\mathbf{P}^1 \arrow[r, "\mathcal{S}_{2,2}\otimes I\otimes I"'] \arrow[d, "\mathcal{S}_{2,\dots,2}"'] & \mathbf{P}^3\times\mathbf{P}^1\times\mathbf{P}^1 . \arrow[d, "I\otimes\mathcal{S}_{2,2}"'] \\
\mathbf{P}^{2^4-1} & \mathbf{P}^3\times\mathbf{P}^3 \arrow[l, "\mathcal{S}_{4,4}"']
\end{tikzcd}$$

For the Segre variety, which is represented by completely decomposable tensor, we have a commuting diagram and $\mathcal{S}_{2,\dots,2} = (\mathcal{S}_{4,4}) \circ (I \otimes \mathcal{S}_{2,2}) \circ (\mathcal{S}_{2,2} \otimes I \otimes I)$.

3 Toric variety and multi-qubit quantum systems

Let $S \subset \mathbf{R}^n$ be finite subset, then a convex polyhedral cone is defined by $\sigma = \mathrm{Cone}(S) = \left\{\sum_{v\in S}\lambda_v v|\lambda_v \geq 0\right\}$. In this case σ is generated by S. In a similar way we define a polytope by $P = \mathrm{Conv}(S) = \left\{\sum_{v\in S}\lambda_v v|\lambda_v \geq 0, \sum_{v\in S}\lambda_v = 1\right\}$. We also could say that P is convex hull of S. A convex polyhedral cone is called simplicial if it is generated by linearly independent set. Now, let $\sigma \subset \mathbf{R}^n$ be a convex polyhedral cone and $\langle u, v\rangle$ be a natural pairing between $u \in \mathbf{R}^n$ and $v \in \mathbf{R}^n$. Then, the dual cone of the σ is define by

$$\sigma^{\wedge} = \{u \in \mathbf{R}^{n*} | \langle u, v\rangle \geq 0 \ \forall\ v \in \sigma\},$$

where $\mathbf{R}^{n*}$ is dual of $\mathbf{R}^n$. We call a convex polyhedral cone strongly convex if $\sigma \cap (-\sigma) = \{0\}$.

The algebra of Laurent polynomials is defined by $\mathbf{C}[z, z^{-1}] = \mathbf{C}[z_1, z_1^{-1}, \dots, z_n, z_n^{-1}]$, where $z_i = \chi^{e_i^*}$. The terms of the form $\lambda \cdot z^\beta = \lambda z_1^{\beta_1} z_2^{\beta_2} \cdots z_n^{\beta_n}$ for $\beta = (\beta_1, \beta_2, \dots, \beta_n) \in \mathbf{Z}$ and $\lambda \in \mathbf{C}^*$ are called Laurent monomials. A ring R of Laurent polynomials is called a monomial algebra if it is a $\mathbf{C}$-algebra generated by Laurent monomials. Moreover, for a lattice cone σ, the ring

$$R_\sigma = \{f \in \mathbf{C}[z, z^{-1}] : \mathrm{supp}(f) \subset \sigma\}$$

is a finitely generated monomial algebra, where the support of a Laurent polynomial $f=\sum_i \lambda_i z^i$ is defined by

$$\mathrm{supp}(f)=\{i\in\mathbf{Z}^n : \lambda_i\neq 0\}.$$

Now, for a lattice cone σ we can define an affine toric variety to be the maximal spectrum

$$\mathbf{X}_\sigma = \mathrm{Spec}R_\sigma.$$

A toric variety $\mathbf{X}_\Sigma$ associated to a fan Σ is the result of gluing affine varieties $\mathbf{X}_\sigma=\mathrm{Spec}R_\sigma$ for all $\sigma\in\Sigma$ by identifying $\mathbf{X}_\sigma$ with the corresponding Zariski open subset in $\mathbf{X}_{\sigma'}$ if σ is a face of $\sigma^{'}$. That is, first we take the disjoint union of all affine toric varieties $\mathbf{X}_\sigma$ corresponding to the cones of Σ. Then by gluing all these affine toric varieties together we get $\mathbf{X}_\Sigma$.

A compact toric variety $\mathcal{X}_A$ is called projective if there exists an injective morphism

$$\Phi : \mathcal{X}_\Sigma \longrightarrow \mathbf{P}^r$$

of $\mathcal{X}_\Sigma$ into some projective space such that $\Phi(\mathcal{X}_\Sigma)$ is Zariski closed in $\mathbf{P}^r$. A toric variety $\mathcal{X}_\Sigma$ is equivariantly projective if and only if Σ is strongly polytopal. Now, let $\mathcal{X}_\Sigma$ be equivariantly projective and morphism Φ be embedding which is induced by the rational map $\phi : \mathcal{X}_A \longrightarrow \mathbf{P}^r$ defined by $p\mapsto [z^{m_0}, z^{m_1},\ldots,z^{m_r}]$, where $z^{m_l}(p)=p^{m_l}$ in case $p=(p_1,p_2,\ldots p_n)$. Then, the rational map $\Phi(\mathcal{X}_\Sigma)$ is the set of common solutions of finitely many monomial equations

$$z_{i_0}^{\beta_0} z_{i_1}^{\beta_1}\cdots z_{i_s}^{\beta_s} = z_{i_{s+1}}^{\beta_{s+1}} z_{i_{s+2}}^{\beta_{s+2}}\cdots z_{i_r}^{\beta_r} \tag{9}$$

which satisfy the following relationships

$$\beta_0 m_0+\beta_1 m_1+\cdots+\beta_s m_s = \beta_{s+1}m_{s+1}+\beta_{s+2}m_{s+2}+\cdots+\beta_r m_r \tag{10}$$

and

$$\beta_0+\beta_1+\cdots+\beta_s=\beta_{s+1}+\beta_{s+2}+\cdots+\beta_r, \tag{11}$$

for all $\beta_l\in\mathcal{Z}_{\geq 0}$ and $l=0,1,\ldots,r$ [3]. As we have seen for multi-qubit systems the separable states are given by the Segre embedding of $\mathbf{CP}^1\times\mathbf{CP}^1\times\cdots\times\mathbf{CP}^1$. Now, for example, let $z_1=\alpha^1_1/\alpha^1_0, z_2=\alpha^2_1/\alpha^2_0,\ldots,z_m=\alpha^m_1/\alpha^m_0$. Then we can cover $\mathbf{CP}^1\times\mathbf{CP}^1\times\cdots\times\mathbf{CP}^1$ by 2^m charts

$$\begin{aligned}
\mathbf{X}_{\check{\Delta}_1} &= \{(z_1,z_2,\ldots,z_m)\},\\
\mathbf{X}_{\check{\Delta}_2} &= \{(z_1^{-1},z_2,\ldots,z_m)\},\\
&\vdots\\
\mathbf{X}_{\check{\Delta}_{2^m-1}} &= \{(z_1,z_2^{-1},\ldots,z_m^{-1})\},\\
\mathbf{X}_{\check{\Delta}_{2^m}} &= \{(z_1^{-1},z_2^{-1},\ldots,z_m^{-1})\}
\end{aligned}$$

Let us consider the m -hypercube Σ centered at the origin with vertices $(\pm1,\ldots,\pm1)$. This gives the toric variety $\mathcal{X}_\Sigma=\mathbf{CP}^1\times\mathbf{CP}^1\times\cdots\times\mathbf{CP}^1$. Now, the map $\Phi(\mathcal{X}_\Sigma)$ is a set of the common solutions of the following monomial equations

$$x_{i_0}^{\beta_0}x_{i_1}^{\beta_1}\cdots x_{i_{2^{m-1}-1}}^{\beta_{2^{m-1}-1}} = x_{i_{2^{m-1}}}^{\beta_{2^{m-1}}}\cdots x_{i_{2^m-1}}^{\beta_{2^m-1}} \tag{12}$$

that gives quadratic polynomials $\alpha_{k_1k_2\ldots k_m}\alpha_{l_1l_2\ldots l_m} = \alpha_{k_1k_2\ldots l_j\ldots k_m}\alpha_{l_1l_2\ldots k_j\ldots l_m}$ for all $j = 1, 2, \ldots, m$ which coincides with the Segre ideals. Moreover, we have

$$\Phi(\mathcal{X}_\Sigma) = \mathrm{Specm}\mathcal{C}[\alpha_{00\ldots0}, \alpha_{00\ldots1}, \ldots, \alpha_{11\ldots1}]/\mathcal{I}(\mathcal{A}), \tag{13}$$

where $\mathcal{I}(\mathcal{A}) = \langle \alpha_{k_1k_2\ldots k_m}\alpha_{l_1l_2\ldots l_m} - \alpha_{k_1k_2\ldots l_j\ldots k_m}\alpha_{l_1l_2\ldots k_j\ldots l_m}\rangle_{\forall j;k_j,l_j=0,1}$. This toric variety describe the space of separable states in a multi-qubit quantum systems. In summary we have investigated the geometrical structures of quantum multi-qubit states based on the Segre variety toric varieties. We showed that multi-qubit states can be characterized and visualized by embedding of toric variety in a complex projective space. The results are interesting in our voyage to the realm of quantum theory and a better understanding of the nature of multipartite quantum systems.

Acknowledgments: The work was supported by the Swedish Research Council (VR).

References

[1] P. Griffiths and J. Harris, *Principles of algebraic geometry*, Wiley and Sons, New York, 1978.

[2] H. Heydari and G. Björk, J. Phys. A: Math. Gen. **38** 3203-3211 (2005).

[3] G. Ewald, *Combinatorial Convexity and Algebraic Geometry*, Springer, 1995.

[4] H. Heydari, *Mathematical Physics Research Developments*, Edited by M. B. Levy, Nova publisher, (2009) pp. 589-604.